

Decay $\Lambda_b \rightarrow pl\bar{\nu}$ in QCD sum rules

Chao-Shang Huang^a, Cong-Feng Qiao^{b,c}, and Hua-Gang Yan^a

^a *Institute of Theoretical Physics, Academia Sinica
P.O. Box 2735, Beijing 100080, China **

^b *CCAST (World Laboratory) P.O. Box 8730, Beijing 100080, China*

^c *ICTP, P.O. Box 586, 34100 Trieste, Italy*

Abstract

The $\Lambda_b \rightarrow p$ semileptonic decay is analyzed by using QCD sum rules within the framework of heavy quark effective theory. The Isgur-Wise function of $\Lambda_b \rightarrow pl\bar{\nu}$ has been calculated. The decay width is given.

PACS: 12.38.Lg, 12.39.Hg, 13.30.Ce, 14.20.Mr.

Keywords: heavy baryon, weak decay, heavy quark effective theory, QCD sum rule.

*Corresponding address

The semileptonic decays of heavy hadrons have been widely investigated as a testing tool for the Standard Model(SM). They can reveal some important features of the physics of heavy quarks with some uncertainties, which are rooted in the nonperturbative aspects of the strong interactions(QCD). Employing the QCD sum rule method [1], we can find a way to estimate the nonperturbative effects, hence can extract some Cabbibo-Kobayashi-Maskawa(CKM) matrix elements by comparing experiments with theoretical calculations.

For the heavy hadrons containing one single heavy quark, an effective theory of QCD based on the heavy quark symmetry in the heavy quark limit [2], the so-called heavy quark effective theory(HQET), has been proposed [3]. For heavy quark to heavy quark transitions, both the classification of the weak decay form factors of heavy hadrons [4, 5] and the computation of the form factors by using sum rules can be greatly simplified in HQET [6–8]. For the heavy quark to light quark semileptonic decay mode, the QCD sum rule method has yet not been used within the framework of HQET. In this letter we shall do such a study.

For heavy to light transitions, one may make use of the heavy quark symmetry for the heavy quark (b or c). It has been shown that the heavy quark symmetry restricts the number of form factors for the heavy baryonic transition $\Lambda_Q \rightarrow \text{light spin-1/2 baryon}$ to two [5]. That is, in the heavy quark limit, the hadronic matrix element of the transition $\Lambda_b \rightarrow p$ is characterized generally by two form factors F_1 and F_2 ,

$$\langle \Lambda_b(v) | \bar{b} \Gamma u | p(P) \rangle = \bar{u}_{\Lambda_b}(v) \Gamma (F_1 + F_2 \not{v}) u_p(P), \quad (1)$$

where v denotes the four-velocity of Λ_b , P denotes the four-momentum of p , and Γ is an arbitrary Dirac matrix ($\Gamma = \gamma_\mu(1 - \gamma_5)$ for $\Lambda_b \rightarrow pl\bar{\nu}$). Hereafter we focus on the determination of F_1 and F_2 by considering the three-point correlators of baryonic currents

$$\tilde{j}^v = \epsilon^{abc} (q_1^{Ta} C \tilde{\Gamma} \tau q_2^b) h_v^c, j = \epsilon^{abc} (q_1^{Ta} C \Gamma_1 \tau q_2^b) \Gamma_2 q_v^c, \quad (2)$$

where $\tilde{j}^v$ is the HQET baryonic current for the heavy quark h_v , while j is the protonic current. There are two choices both for $\tilde{j}^v$ and j . We choose $\tilde{\Gamma} = \gamma_5$ for $\tilde{j}^v$ for the sake of simplicity since the numerical differences resulting from the different choices of $\tilde{j}^v$ are not significant [8]. We choose the tensor variant for j since it seems to be more suitable for studying the electromagnetic and strong properties of light baryons [9], i.e.

$$\begin{aligned} j &= \epsilon^{abc} u^a C \sigma_{\mu\nu} u^b \sigma^{\mu\nu} \gamma_5 d^c \\ &= 4\epsilon^{abc} (u^a C \gamma_5 d^b u^c + u^a C d^b \gamma_5 u^c). \end{aligned} \quad (3)$$

The three point correlator in present case is

$$\Pi(P', P, z) = i^2 \int d^4x d^4y e^{ik \cdot x - iP \cdot y} \langle 0 | T \tilde{j}^v(x) \bar{h}_v(0) \Gamma u(0) \bar{j}(y) | 0 \rangle, \quad (4)$$

where $P' = m_b v + k$ and $z = P \cdot v$. After inserting a complete set of physical intermediate states, as the phenomenological consequence of (4), we have

$$\Pi(P', P, z) = f_{\Lambda_b} f_p \frac{2}{(\omega - 2\bar{\Lambda})} P_+ \Gamma [F_1(z) + F_2(z) \not{v}] \frac{\not{P} + m_p}{P^2 - m_p^2} + \text{res}, \quad (5)$$

where $P_+ = (1 + \not{v})/2$, $\bar{\Lambda} = m_{\Lambda_b} - m_b$, $\omega = 2k \cdot v$ and f_{Λ_b} , f_p are the so-called "decay constants" which are given by

$$\langle 0 | \tilde{j}^v | \Lambda_b \rangle = f_{\Lambda_b} u, \quad \langle p(P) | \bar{j} | 0 \rangle = f_p \bar{u}(P). \quad (6)$$

They can be found in Ref. [8] and Refs. [10, 11], respectively. To obtain (5) we have taken into account (1) and the heavy quark limit.

Introducing the assumption of quark-hadron duality, the contribution from higher resonant and continuum states can be treated as

$$res. = \int_{D'} d\nu ds \frac{\rho_{pert}(\nu, s, z)}{(\nu - \omega)(s - P^2)}. \quad (7)$$

The region D' is characterized by one or two continuum thresholds ν_c, s_c . As a essential practice in QCD sum rule, in order to incorporate the assumption, we should express the perturbative term in the form of dispersion relation

$$\Pi_{pert}^i(\omega, P^2, z) = \int d\nu ds \frac{\rho^i(\nu, s, z)}{(\nu - \omega)(s - P^2)}, \quad (i = 1, 2), \quad (8)$$

where $i = 1, 2$ denote the different terms associated with F_1 and F_2 , respectively.

The calculation of (4) can be done by using operator product expansion(OPE)

$$T[\tilde{j}^v(x)\bar{h}_v(0)\Gamma u(0)\bar{j}(y)] = \Sigma_n C_n(x, y)O_n(0). \quad (9)$$

The first term represents the perturbative contribution, while the remaining terms represent the nonperturbative effects after introducing the QCD vacuum and condensates. Up to dimension 6, the relevant Feynman diagrams to compute (4) are plotted in Fig.1.

In our calculation, the coordinate representation is adopted. The heavy quark propagator

$$\langle Th_v(x)\bar{h}_v(0) \rangle = \int_0^\infty dt \delta(x - tv) P_+$$

and the fixed point gauge [12] are used. For our purposes it is sufficient to retain the condensates with dimensions lower than 7. We use the following values of the condensates [13]:

$$\begin{aligned} \langle \bar{q}q \rangle &\simeq -(0.23 \text{ GeV})^3, \\ \langle \alpha_s GG \rangle &\simeq 0.04 \text{ GeV}^4. \end{aligned} \quad (10)$$

The normalization $\text{Tr}\tau^\dagger\tau = 1$ has been used in the analysis. In the fixed-point gauge, the space-time translational invariance is violated, but it is restored by adding all the diagrams in Fig. 1. This is a check of our calculation.

In the standard way, we employ a double Borel Transformation $\omega \rightarrow M, P^2 \rightarrow T$ in order to suppress the higher excited states and continuum states contributions. The analytic expressions we got for F_1 and F_2 after Borel Transformation are:

$$\begin{aligned} -2f_{\Lambda_b}f_pF_1e^{-2\bar{\Lambda}/M-m_p^2/T} &= \int_0^{\nu_c} d\nu \int_{m_p^2}^{2\nu z} ds \rho_{pert}^1 e^{-s/T-\nu/M} - \frac{1}{3}\langle \bar{q}q \rangle^2 - \\ &\quad \frac{1}{32\pi^4}\langle \alpha_s GG \rangle \int_0^{T/4} \left(1 - \frac{4\beta}{T}\right) e^{-4\beta(1-4\beta/T)/M^2-8\beta z/(TM)} d\beta, \\ -2f_{\Lambda_b}f_p m_p F_2 e^{-2\bar{\Lambda}/M-m_p^2/T} &= \int_0^{\nu_c} d\nu \int_{m_p^2}^{2\nu z} ds \rho_{pert}^2 e^{-s/T-\nu/M} + \\ &\quad \frac{1}{8\pi^4}\langle \alpha_s GG \rangle \int_0^{T/4} \left(1 - \frac{4\beta}{T}\right) \frac{\beta}{M} e^{-4\beta(1-4\beta/T)/M^2-8\beta z/(TM)} d\beta, \end{aligned} \quad (11)$$

where

$$\begin{aligned} \rho_{pert}^1 &= \frac{1}{32\pi^4\sigma^3} [-2z^3\sigma^3 - (-s + z(\nu + 2z))^3 + 3z^2(-s + z(\nu + 2z))\sigma^2], \\ \rho_{pert}^2 &= \frac{1}{64\pi^4\sigma^3} [s - 2z^2 + z(-\nu + \sigma)]^2 [\nu s + 8z^3 - 4z^2(-2\nu + \sigma) - 2z(-\nu^2 + 5s + \nu\sigma)], \end{aligned} \quad (12)$$

with $\sigma = \sqrt{-4s + (\nu + 2z)^2}$. One can see from (11) that the four-quark condensate represents the main nonperturbative contribution to F_1 , which is similar to the case of heavy to heavy transitions [14].

In our numerical analysis, the "decay constants" and some other constants we used are [8, 10, 15]:

$$\begin{aligned} m_{\Lambda_b} &= 5.64\text{GeV}, m_p = 0.938\text{GeV}, f_{\Lambda_b} = \sqrt{0.0003}\text{GeV}^3 \\ f_p &= 0.0255\text{GeV}^3, \Lambda = 0.79\text{GeV}. \end{aligned} \quad (13)$$

It should be mentioned here that the values of f_p defined in (6) is probably different from which we cited from Ref. [10] because we have adopted a different proton current, that is, the "tensor current", instead of the "vector current" [10, 11]. However, if the approximations in sum rule calculations are justified to be good enough, these two currents should give roughly the same value of f_p . We find that with the threshold $\nu_c = 3.7\text{GeV}$, we can have a reasonably good window for F_1 , where $1.4\text{GeV} < \frac{4T}{m_b} = M < 1.8\text{GeV}$. The results are given in Fig.2 and Fig.3 respectively, where the different curves correspond to different choices of the Borel parameters.

The semileptonic decay $\Lambda_b \rightarrow pl\bar{\nu}$ can be analyzed directly after we obtain the Isgur-Wise function F_1 and F_2 . By neglecting the lepton mass (for $l=e, \mu$), it is easy to show that the differential decay width is:

$$\begin{aligned} \frac{d\Gamma}{dz} &= \frac{|V_{ub}|^2 G_F^2}{12\pi^3} [F_1^2(3m_p^2 z - 2m_p^2 m_{\Lambda_b} + 3m_{\Lambda_b}^2 z - 4m_{\Lambda_b} z^2) + 2F_1 F_2(m_p^3 + 3m_p m_{\Lambda_b}^2 - 6m_{\Lambda_b} m_p z \\ &\quad + 2m_p z^2) + F_2^2(2m_p^2 m_{\Lambda_b} - m_p^2 z + 3m_{\Lambda_b}^2 z - 8m_{\Lambda_b} z^2 + 4z^3)] \sqrt{z^2 - m_p^2}. \end{aligned} \quad (14)$$

The numerical results of $\frac{d\Gamma}{dz}$ are shown in Fig.4. The total width are $\Gamma = 1.35 \times 10^{-11} |V_{ub}|^2 \text{GeV}$. The comparing values are $\Gamma = 1.50 \times 10^{-11} |V_{ub}|^2 \text{GeV}$, when $M = 1.4\text{GeV}$; $\Gamma = 1.40 \times 10^{-11} |V_{ub}|^2$, when $M = 1.8\text{GeV}$.

There are a few comments to be made regarding the results and our computation:

(a) The choice of $\frac{4T}{m_b} = M$ is purely for convenience. We find that the $\max(T, M)$ matters in the computation when the magnitudes of T and M are diverse greatly. The choice is somewhat similar to that of Ref. [16], where the analyze of semileptonic decay of B meson was performed. The values of Borel parameters we used seem very large compared with that of Ref. [14]. In fact, we encountered the same situation mentioned in Ref. [17], i. e., the influence of the four-quark condensate is still large for the values of Borel parameters. Nevertheless, comparing to those in Ref. [16], the values are similar. It seems to be natural that the Borel parameters in heavy to light transitions may take values different from those in heavy to heavy transitions.

(b) Similar to the case in Ref. [18], we find that F_1 is dominated by the four-quark condensate (about 63%) instead of the perturbative term. Therefore, to assume the hybrid sum rules [18] may be better in order to calculate F_1 . As for F_2 , because it lacks the four-quark condensate, the perturbative term dominates. F_2 has no good stability in the window. This is because the α_s corrections which are expected to be more important for F_2 than for F_1 have not been included in this paper. However, we could obtain a good window for $\frac{d\Gamma}{dz}$ and for the total width.

(c) The absolute values of F_1 and F_2 depend on the "decay" constants of Λ_b and p , which themselves have some uncertainties. The result for decay width will double these uncertainties. Furthermore, there exists the uncertain CKM element $|V_{ub}|$ in the theoretical determination of the magnitude of decay width. In order to eliminate these uncertainties one considers the ratio $R = F_2/F_1$. Our numerical result for $R \simeq -0.42$ at zero recoil ($z = m_p$)

is in consistence with the experimental data for $\Lambda_b \rightarrow \Lambda e \bar{\nu}$ [19]. This is expected in both the heavy quark limit and the light flavor SU_3 limit. Moreover, our result suggests a tendency of the growth of R with the final state baryon getting lighter. Note that if the final state baryon is a heavy baryon, R will approach zero. So the tendency is in the right direction.

In summary, we have calculated the form factors of $\Lambda_b \rightarrow p$ in the $m_b \rightarrow \infty$ limit from QCD sum rules within the framework of HQET. We have also calculated the decay width of $\Lambda_b \rightarrow pl\bar{\nu}$ using the obtained form factors. With minor modifications, the results can be generalized to $\Lambda_c \rightarrow \Lambda l\bar{\nu}$ which we shall analyze in detail elsewhere.

We thank Y. B. Dai for discussions. C.F.Qiao wishes to thank the ICTP high energy physics group for the hospitality during his visit. This work is supported in part by the National Natural Science Foundation of China and the Hua Run Postdoctoral Science Foundation of China.

REFERENCES

- [1] M.A. Shifman, A.I. Vainshtein and V.I. Zakharov, Nucl. Phys. B 147 (1979)385, 488.
- [2] N. Isgur and M.B. Wise, Phys. Lett. B 232 (1989) 113, B 237 (1990) 527;
M.B. Voloshin and M.A. Shifman, Yad. Fiz. 45 (1987) 463, 47 (1988) 801.
- [3] H. Georgi, Phys. Lett. B 240 (1990) 447;
A.F. Falk, H. Georgi, B.Grinstein and M.B. Wise, Nucl. Phys. B343 (1990) 1.
- [4] N. Isgur and M.B. Wise, Nucl. Phys. B 348 (1991) 276;
H. Georgi, Nucl. Phys. B 348 (1991) 293;
W. Roberts, Nucl. Phys. B 389 (1993)549;
Y.B. Dai, X.H. Guo and C.S. Huang, Nucl. Phys. B412 (1994) 277.
- [5] T. Mannel, W. Roberts and Z. Ryzak, Nucl. Phys. B 355 (1991) 38.
- [6] P. Ball, Phys. Rev. D48 (1993) 3190;
A.G. Grozin and O.I. Yakovlev, Phys. Lett. B 285(1992) 254, B 291 (1992) 441;
Y.B. Dai, C.S. Huang and M.Q. Huang, Phys. Rev. D 55 (1997) 5719.
- [7] M. Neubert, Phys. Rev. D46 (1992) 1076, 3914.
- [8] Y.B. Dai, C.S. Huang, C. Liu and C.D. Lü, Phys. Lett. B 371 (1996) 99.
- [9] G.V. Etimov, M.A. Ivanov and V.E. Lyubovitskij, Z. Phys. C 47 (1990) 583.
- [10] V.A. Nesterenko and A. V. Radyushkin, Phys. Lett. B 128 (1983) 439.
- [11] E.V. Shuryak, Nucl. Phys. B 198 (1982) 83.
- [12] For a review, see V.A. Novikov, M.A. Shifman, A.I. Vainshtein and V.I. Zakharov, Fortschr. Phys. 32 (1984) 585.
- [13] B.L. Ioffe and A.V. Smilga, Phys. Lett. B114 (1982) 353; V.A. Nesterenko A.V. Radyushkin, Phys. Lett. B115 (1982) 410.
- [14] Y.B. Dai, C.S. Huang, M.Q. Huang and C. Liu. Phys. Lett. B 387 (1996) 379.
- [15] Particle Data Group, Phys. Rev. D54 (1996) 1.
- [16] P. Ball, Phys. Rev. D48 (1993) 3190.
- [17] H.G. Dosch, E. Ferreira, M. Nielsen and R. Rosenfeld, hep-ph/9712350.
- [18] S. Narison, Phys. Lett. B 345 (1995) 166.
- [19] CLEO Collaboration, G. Crawford *et al.*, Phys Rev. Lett. 75, (1995) 624; CLEO Collaboration, T. Bergfeld *et al.*, Report No. CLEO 94-4,1994 (unpublished).

Figure Captions

Fig. 1 Feynman diagrams for the computation of $\Pi(P', P, z)$. The diagrams where the gluon condensates are generated from b -quark line are not plotted because the contributions corresponding to the diagrams are zero due to the use of fix-point gauge.

Fig. 2 Isgur-Wise function $F_1(z)$ with different values of Borel parameter T and M.

Fig. 3 Isgur-Wise function $F_2(z)$ with different values of Borel parameter T and M.

Fig. 4 The differential decay width with different values of Borel parameter.




Fig. 4




Fig. 2




Fig. 3




Fig. 1